

# Immunohistochemical Distribution of Type IV Collagenase in Normal, Benign, and Malignant Breast Tissue

Carlos Monteagudo, Maria J. Merino, Josefina San-Juan, Lance A. Liotta, and William G. Stetler-Stevenson

From the Laboratory of Pathology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

*Production of type IV collagenase by tumor cells has been linked to their metastatic potential in several experimental models. A possible role for this enzyme in basement membrane type IV collagen turnover has also been suggested. Two recently developed affinity-purified, monospecific antibodies directed against the amino terminus (H<sub>1</sub>), or an internal active site domain (metal binding region [MBR]) of human type IV collagenase, were employed in the avidin-biotin-immunoperoxidase technique in formalin-fixed, paraffin-embedded breast tissue samples from 55 patients. Intense cytoplasmic immunostaining of myoepithelial cells was found in normal and hyperplastic tissue, and discontinuous staining was noted in intraductal carcinomas. Luminal epithelial cells were negative or weakly positive in large- or medium-sized ducts but reacted frequently in normal terminal ducts and hyperplastic lesions. Epithelial cells in intraductal carcinomas exhibited immunoreactivity in 20 of 23 cases. Invasive carcinomas were positive in 36 of 40 cases, and metastatic cells in lymph nodes stained in 10 of 12 cases. These results support a role for type IV collagenase in the basement membrane remodeling of normal breast. Our findings suggest that myoepithelial cells play a pivotal role in this enzymatic activity. The high percentage of positive cells in invasive carcinomas and the strong immunoreactivity of lymph node metastases support the role of the enzyme in tumor invasion and metastasis and suggest that tumor cells are the essential source of the enzyme in these processes. (Am J Pathol 1990, 136:585-592)*

During the last decade, increasing attention has been focused on the role of extracellular matrix, especially basement membrane (BM), in the process of tumor invasion and metastasis.<sup>1-7</sup> Although the biochemical mechanisms involved in BM turnover under physiologic conditions remain essentially unknown,<sup>8</sup> it has been established that the BM-degrading properties of tumor cells correlate with their metastatic potential.<sup>9</sup>

Type IV collagen is one of the major components of BM, and it represents the structural scaffolding of these specialized sheets of extracellular matrix. The enzymatic degradation of type IV collagen is specifically initiated by a neutral metalloproteinase, type IV collagenase.<sup>10,11</sup> This enzyme has been found in human tumor cells<sup>12,13</sup> as well as in other normal cell types, such as endothelium,<sup>14</sup> fibroblasts,<sup>13</sup> macrophages,<sup>15</sup> polymorphonuclear leukocytes,<sup>16</sup> and keratinocytes.<sup>17</sup> It is secreted in a latent form that can be activated, at least *in vitro*, by trypsin and organomercurial compounds. This activation is associated with the loss of 80 amino acid residues from the amino terminus.<sup>18</sup>

The structure and changes of BM in breast lesions have been extensively studied,<sup>19-24</sup> and a variable loss of BM is a constant feature in invasive neoplasms. Previous investigators, employing polyclonal antibodies against murine type IV collagenase, have found positive immunoreactivity in invading breast cancer cells but not in normal breast, benign lesions, or *in situ* carcinomas.<sup>25</sup>

We have used two recently developed affinity-purified rabbit polyclonal antibodies directed against synthetic peptides corresponding to different domains of the human type IV collagenase in an attempt to study the immunohistochemical distribution of this enzyme in normal and hyperplastic breast tissue as well as in primary malignant lesions and metastases.

## Materials and Methods

### Tissues

Breast tissue samples from 55 patients were obtained from the files of the Laboratory of Pathology, National

---

Accepted for publication October 20, 1989.

Address reprint requests to Maria J. Merino, MD, Laboratory of Pathology, National Cancer Institute, Bldg. 10, Room 2N212, Bethesda, MD 20892.

Cancer Institute. The tissues were routinely fixed in 10% neutral buffered formalin and embedded in paraffin. Non-tumorous tissue included 14 cases of normal breast, six cases of fibrocystic changes, six fibroadenomas, 11 breast lesions showing epithelial hyperplasia (four mild, three moderate, and four florid), and one case of sclerosing adenosis. Forty carcinomas were evaluated: 37 ductal type (34 not otherwise specified, [NOS], two tubular, and one with squamous differentiation) and three lobular. Four cases were classified as pure intraductal carcinoma, but *in situ* tumors were found as a component of 19 invasive cancers. Lobular carcinoma *in situ* was present in all three cases of infiltrating lobular carcinoma. Twelve axillary lymph node metastases from five patients were included in the study.


Frozen tissue samples were also evaluated to compare the reliability of the immunohistochemical results with that of formalin-fixed material.

### Antibody Characterization

Anti-type IV collagenase antibodies were prepared with the use of synthetic peptides coupled to bovine serum albumin. Peptide H<sub>1</sub> with the sequence APSIIKFPGD-VAPKTD, corresponding to the 17 amino-terminal amino acid residues of type IV procollagenase, and peptide MBR with the sequence VAAHEFGHAMGLEHSQ, corresponding to the putative metal ion-binding domain of type IV collagenase, were synthesized on a Bioscience 9600 peptide synthesizer (Bioscience, Novato, CA) with tBOC amino acid methodology. The peptides were coupled to bovine serum albumin with glutaraldehyde (0.14%). For the initial immunizations, 1 ml of bovine serum albumin-peptide conjugate was mixed with 1 ml of complete Freund's adjuvant and emulsified prior to subcutaneous injection. For the remaining biweekly immunization, 0.5 ml of bovine serum albumin-peptide conjugate was emulsified with 0.5 ml of incomplete Freund's adjuvant before injection.

Peptide affinity resins were prepared for antibody purification for both antibodies with Affi-Gel 10 (BioRad, Richmond, CA), following the manufacturer's directions and using 2 mg of each peptide. These resins were used to affinity purify the antibodies from rabbit serum, following heat inactivation of the serum at 56°C for 30 minutes. Following absorption of the antibodies onto the resin overnight at 4°C, the columns were washed with 20 column volumes of cold phosphate-buffered saline prior to elution with two bed volumes of 1 mol acetic acid. This eluate was immediately neutralized by the addition of 1 mol NaOH before buffer exchange to phosphate-buffered saline using a YM 10 membrane (Amicon, Danvers, MA).

Purified antibodies H<sub>1</sub> and MBR were stored at 4°C and characterized with Western blot analysis or enzyme-



**Figure 1.** Immunoblotting of purified type IV collagenase and conditioned media samples using anti-type IV collagenase synthetic peptide antibodies. All samples were run on a 10% polyacrylamide gel using a Laemmli buffer system and electroblotted to Immobilon membranes using a Tris-glycine/methanol transfer buffer system. A: 2 µg of gelatin-affinity purified type IV collagenase. Primary antibody H<sub>1</sub>. B: 10 µl of human melanoma cell conditioned media containing approximately 80 ng of type IV collagenase. Primary antibody H<sub>1</sub>. The single band indicates that other neutral metalloproteinases present in this sample do not cross react with the primary antibody. C: 10 µl of human melanoma cell conditioned media containing approximately 80 ng of type IV collagenase. Primary antibody MBR. The single band indicates that other neutral metalloproteinases present in this sample do not cross react with the primary antibody.

linked immunosorbent assay (ELISA) before use in the immunohistochemical studies. The results of Western blot analyses showed that both antibody preparations were monospecific and reacted with a single species in human melanoma cell (A2058) conditioned media that corresponds to purified type IV collagenase (Figure 1).

### Immunohistochemical Procedure

The avidin-biotin-immunoperoxidase technique was used as previously described by Hsu et al.<sup>26</sup> After the sections were dewaxed and rehydrated, endogenous peroxidase activity was blocked by incubating the slides in 0.3% H<sub>2</sub>O<sub>2</sub> in absolute methanol for 30 minutes. The tissue samples were then exposed for 20 minutes to 2% normal goat serum. Without washing, first antibodies (H<sub>1</sub> or MBR) were incubated for 45 minutes in a humidity chamber at room temperature. Dilutions were 1 to 2 µg/ml for H<sub>1</sub> and

**Table 1.** Distribution of Type IV Collagenase Immunoreactivity in Normal Breast and Benign Breast Lesions

Tissue*	Myoepithelium†	Luminal/ ductal epithelium‡
Normal breast (14)	+++	-/+‡
Benign lesions		
Fibrocystic condition (6)	++	-/+
Epithelial hyperplasia (11)	++	+/>++§
Sclerosing adenosis (1)	++	-/+
Fibroadenoma (6)	variable	variable

\* Number of cases in parentheses.

† -, no staining; -/+, occasional weak staining; +, weak staining; ++, moderate staining; +++, intense staining.

‡ Mainly in terminal ducts.

§++ in florid hyperplasia.

15 µg/ml for MBR. Second biotinylated goat anti-rabbit IgG (dilution 1/200) was incubated for 30 minutes. The slides were then incubated in avidin-biotin-peroxidase (ABP) complex (Vectastain kit, Vector Laboratories, Burlingame, CA) for 45 minutes, and the reaction was revealed by 0.5 mg/ml 3,3'-diaminobenzidine tetrahydrochloride (Sigma, St. Louis, MO) and 3 µl/ml H<sub>2</sub>O<sub>2</sub> in 50 mmol TRIS, pH 7.6. All antibodies were diluted in 50 mmol TRIS-buffered saline, pH 7.5. Between steps, the slides were washed three times in TRIS-buffered saline. Control slides were performed by replacing first antibody with the IgG fraction of normal rabbit serum. The sections were considered as positive or negative according to the presence or absence of specific staining and were evaluated on an Olympus microscope (Optical Elements Corp., Washington, DC). The terms "terminal duct" and "ductule" are used here as described by Wellings et al.<sup>27</sup>

## Results

The pattern of staining was similar with both antibodies; therefore, we will consider them together except for those cases in which marked differences were noted (Tables 1 and 2). Immunoreactivity appeared as a cytoplasmic, granular, or diffuse staining in all cases. Formalin-fixed as well as frozen tissue showed areas without immunostaining.

### Normal Breast

Strong immunostaining of myoepithelial cells (Figure 2A) was found in normal ducts and ductules. This reactivity occurred with both antibodies, but it was more intense with H<sub>1</sub>. In areas with involuting changes, such as myoid atrophy, myoepithelium was also strongly stained. Luminal epithelial cells were negative or weakly positive in

large- or medium-sized ducts, but in terminal ducts they occasionally showed an enhanced immunoreactivity. In one case of lactating breast associated with an invasive carcinoma, myoepithelial cells were strongly positive, while secretory epithelium was completely negative (Figure 3).

### Benign Lesions

When fibrocystic changes were noted, the cysts covered by apocrine metaplasia were occasionally positive, as were some cysts with flattened epithelium. Hyperplastic lesions revealed an increased reactivity of the proliferative epithelium, while myoepithelial cells also remained positive. The immunostaining was usually stronger in cases of florid hyperplasia. Sclerosing adenosis showed a positive reaction in most cells with H<sub>1</sub>. The results in fibroadenomas were completely variable, being negative in some cases and showing indistinct staining of myoepithelium, epithelium, or both in the remaining cases. Similar results were found in frozen tissue.





### In Situ Carcinomas

Intraductal carcinomas with adjacent invasive component showed positive immunostaining in 16 cases (84%) in at least 10% of the tumor cells (Figures 2B and 4). The epithelial cells in intraductal carcinomas reacted markedly stronger than did those of hyperplastic lesions. A positive, discontinuous, myoepithelial cell layer was present in most cases. Four of four cases of ductal cancer without invasion were positive, and there was no correlation with histologic type (comedo vs. noncomedo). Three of three *in situ* lobular carcinomas, all of them associated with invasive lobular carcinomas, were also immunoreactive.

**Table 2.** Frequency of Intense Epithelial Cell Immunostaining with Anti-type IV Collagenase Antibodies in Breast Cancer Tissues

Diagnosis	No. of positive cases/ total no. cases	Percentage
I. <i>In situ</i> carcinomas		
Intraductal		
with invasion	16/20	84
without invasion	4/4	100
Lobular, associated		
with invasion	3/3	100
II. Invasive carcinomas		
Ductal	33/37	89
Lobular	3/3	100
III. Metastases	10/12	83







## Invasive Carcinomas

Infiltrating ductal carcinomas were positive in 33 cases (89%) in approximately 80% of the tumor cells (Figure 2C). Invasive lobular carcinomas were also positive in the three cases studied.

## Metastases

Metastatic tumor cells in lymph nodes showed immunoreactivity in 10 cases (83%) (Figure 2D). Lymphoid cells were negative, but histiocytes showed occasional positivity of the cytoplasm.

In all samples, smooth muscle cells were immunoreactive. Endothelial cells rarely exhibited a positive staining, mainly in areas of vascular neogenesis. Fibroblasts were mostly negative, although a few immunoreactive ones with myofibroblastlike appearance were found. The frozen tissues stained in a pattern similar to that of the formalin-fixed tissues.

## Discussion

The cellular source of extracellular matrix-degrading enzymes has important implications in our understanding of tumor biology and tissue remodeling.<sup>7</sup> During invasion and the process of metastases, tumor cells must traverse epithelial and endothelial BM, where type IV collagen represents a barrier as a major structural component.<sup>9</sup>

Host-tumor cell interactions are known to play a role in the degradation of interstitial (I, II, and III) collagens.<sup>4,28</sup> Pepsinized type IV collagen is not degraded by classical interstitial collagenases,<sup>29</sup> although it is sensitive to some extent to other nonspecific proteases such as elastase, cathepsin G, and plasmin.<sup>3</sup> The isolation and characterization of type IV collagenase<sup>10,11</sup> introduced a new approach in BM physiology and pathology. The role of this enzyme in tumor invasion has been discussed previously.<sup>1,2,6,30</sup> Experimental evidence of a link between the increased production of type IV collagenase by tumor cells and metastatic behavior was described in murine models,<sup>9</sup> and recently it has been extended to the *ras*-induced metastatic phenotype.<sup>31</sup>

Our findings of a marked increase in type IV collagenase immunoreactivity in invasive breast carcinomas are in accordance with previous studies.<sup>25</sup> Reactivity in *in situ*

carcinomas and in lymph node metastases, however, has not been previously documented. Immunostaining in *in situ* carcinomas is a common finding, and it does not seem to be related to the histologic type (comedo vs. non-comedo). Differences between cases with and without invasive components are not significant in our material, but because of the small number of cases without invasion, additional studies are required to complement our data.



Whether *in situ* or invasive, it seems that tumor cells are the essential source of this enzyme, with no major detectable participation of fibroblasts or other cell types in the tumor stroma. This directly supports the concept proposed by Liotta and coworkers<sup>9,25</sup> that increased expression of type IV collagenase is a marker of malignant conversion. Also, this observation contrasts with the known production of interstitial collagenases by normal fibroblasts, near the invasion front of human basal cell carcinomas,<sup>32,33</sup> or in culture, stimulated by a membrane-associated factor from tumor cells.<sup>28,34</sup> The latter investigators, however, were not able to find production of type IV collagenase by their cultured fibroblasts.<sup>28</sup>

The increased immunoreactivity of epithelial cells in cases of epithelial hyperplasia was less intense than that in *in situ* carcinomas. The interpretation of this observation awaits further analysis. In regard to the inconstant pattern of reactivity in fibroadenomas, some authors have found similar results in these lesions when using anti-actin antibodies as myoepithelial cell markers.<sup>24</sup>

Martinez-Hernandez et al<sup>35</sup> found that an unidentified enzymatic activity was involved in BM collagen turnover in involuting mammary glands. Liotta and coworkers<sup>10,36</sup> have suggested that type IV collagenase could be involved in the breast BM remodeling. Our results of type IV collagenase immunoreactivity in normal breast support the role of the enzyme in these normal processes. Myoepithelial cells seem to be the main cell type involved in this enzymatic activity, although the participation of luminal epithelial cells is increased in terminal ducts.

Although the best-known function of myoepithelium is its contractile ability, a possible role of these cells in BM formation was initially suggested<sup>37,38</sup> and later confirmed *in vitro*.<sup>39,40</sup> Warburton et al<sup>41</sup> found intracellular immunostaining with anti-laminin and type IV collagen antibodies in the basal (myoepithelial-like) cells of the terminal end buds of 7-day-old rats. These authors considered that these cells were actively synthesizing BM components. Now we support the role for myoepithelium in BM collagen turnover, based on the immunoreactivity of this cell type

**Figure 2.** Type IV collagenase immunoreactivity in normal and neoplastic breast tissue. **A (Top left):** Normal breast. Terminal ducts showing strong immunostaining of myoepithelial cells. (Imm. H<sub>1</sub>, 250X) **B (Top right):** Intraductal carcinoma. The epithelial cells were markedly positive. (Imm. H<sub>1</sub>, 150X) **C (Bottom left):** Invasive ductal carcinoma. Notice the intense cytoplasmic immunostaining of the tumor cells. (Imm. MBR, 250X) **D (Bottom right):** Lymph node. Metastatic tumor cells are positive, while the lymphocytes are unreactive. (Imm. MBR, 250X).



**Figure 3.** Lactating breast. The myoepithelial cells show strong positivity. The secretory epithelial cells are negative (Imm. H<sub>1</sub>, 250X).

**Figure 4.** Heterogeneous type IV collagenase immunoreactivity in intraductal carcinoma. In this case, the majority of the cells show negative or weakly positive staining (Imm. H<sub>1</sub>, 250X).

for type IV collagenase. The immunostaining of luminal epithelial cells in terminal ducts is not surprising because these cells are also reactive with a myoepithelial cell marker such as KA 1 antibody.<sup>42,43</sup>

In summary, we have used affinity-purified, monospecific antibodies against human type IV collagenase to study the distribution of this enzyme in normal and diseased breast. The results of our study demonstrate that in normal breast tissue type IV collagenase is primarily confined to the myoepithelial cells. We propose that this distribution reflects the participation of these cells in the physiologic turnover of the basement membrane that surrounds breast ducts and ductules. Furthermore, we demonstrate an intense epithelial immunoreactivity with these antibodies in intraductal carcinomas (with and without invasive carcinoma), lobular carcinoma *in situ* (associated with adjacent invasive carcinoma), and invasive carcinomas (ductal and lobular), as well as associated lymph node metastases. Our results clearly show a redistribution in type IV collagenase immunoreactivity from predominantly myoepithelial association in normal and benign breast tissue to an intense epithelial cell staining in malignant breast lesions. This suggests that these antibodies may serve as a useful tool for the evaluation of malignant breast disease.

## References

- Liotta LA, Rao CN, Barsky SH: Tumor invasion and the extracellular matrix. *Lab Invest* 1983, 49:636-649
- Liotta LA: Tumor invasion and metastases: Role of the basement membrane. *Am J Pathol* 1984, 117:339-348
- Pauli BU, Kuettner KE: Tumor invasion and its local regulation. *Urology* 1984, 23(Suppl):18-28
- Woolley DE: Collagenolytic mechanisms in tumor cell invasion. *Cancer Metastasis Rev* 1984, 3:361-372
- Albrechtsen R, Wewer UM, Liotta LA: Basement membranes in human cancer. *Pathol Annu* 1986, 21(2):251-276
- Liotta LA: Tumor invasion and metastases—Role of the extracellular matrix: Rhoads Memorial Award lecture. *Cancer Res* 1986, 46:1-7
- Pauli BU, Knudson W: Tumor invasion: A consequence of destructive and compositional matrix alterations. *Hum Pathol* 1988, 19:628-639
- Timpl R, Dziadek M: Structure, development and molecular pathology of basement membranes. *Int Rev Exp Pathol* 1986, 29:1-112
- Liotta LA, Tryggvason K, Garbisa S, Hart I, Foltz CM, Shafie S: Metastatic potential correlates with enzymatic degradation of basement membrane collagen. *Nature* 1980, 284:67-68
- Liotta LA, Abe S, Robey PG, Martin GR: Preferential digestion of basement membrane collagen by an enzyme derived from a metastatic murine tumor. *Proc Natl Acad Sci USA* 1979, 76:2268-2272
- Liotta LA, Tryggvason K, Garbisa S, Robey PG, Abe S: Partial purification and characterization of a neutral protease which cleaves type IV collagen. *Biochemistry* 1981, 20:100-108
- Salo T, Liotta LA, Keski J, Turpeenniemi-Hujanen T, Tryggvason K: Secretion of basement membrane collagen degrading enzyme and plasminogen activator by transformed cells—Role in metastasis. *Int J Cancer* 1982, 30:669-673
- Collier IE, Wilhelm SM, Eisen AZ, Marmar BL, Grant GA, Seltzer JL, Kronberger A, He C, Bauer EA, Goldberg GI: H-ras oncogene-transformed human bronchial epithelial cells (TBE-1) secrete a single metalloprotease capable of degrading basement membrane collagen. *J Biol Chem* 1988, 263:6579-6587
- Kalebic T, Garbisa S, Glaser B, Liotta LA: Basement membrane collagen: Degradation by migrating endothelial cells. *Science* 1983, 221:281-283
- Garbisa S, Ballin M, Daga-Giordini D, Fastely G, Naturale M, Negro A, Semenzato G, Liotta LA: Transient expression of type IV collagenolytic metalloproteinase by human mononuclear phagocytes. *J Biol Chem* 1986, 261:2369-2375
- Uitto VJ, Schwartz D, Veis A: Degradation of basement membrane collagen by neutral proteases from human granulocytes. *Eur J Biochem* 1980, 105:409-417
- Woodley DT, Kalebic T, Banes AJ, Link W, Prunieras M, Liotta LA: Adult human keratinocytes migrating over nonviable dermal collagen produce collagenolytic enzymes that degrade type I and type IV collagen. *J Invest Dermatol* 1986, 86:418-423
- Stetler-Stevenson WG, Kruttsch HC, Wachter MP, Margulies IMK, Liotta LA: The activation of human type IV collagenase proenzyme. Sequence identification of the major conversion product following organomercurial activation. *J Biol Chem* 1989, 264:1353-1356
- Siegal GP, Barsky SH, Terranova VP, Liotta LA: Stages of neoplastic transformation of human breast tissue as monitored by dissolution of basement membrane components. *Invasion Metastasis* 1981, 1:54-65
- Albrechtsen R, Nielsen M, Wewer U, Engvall E, Ruoslahti E: Basement membrane changes in breast cancer detected by immunohistochemical staining for laminin. *Cancer Res* 1981, 41:5076-5081
- Gusterson BA, Warburton MJ, Mitchell D, Ellison M, Neville AM, Rudland PS: Distribution of myoepithelial cells and basement membrane proteins in the normal breast and in benign and malignant breast diseases. *Cancer Res* 1982, 42:4763-4770
- Barsky SH, Siegal G, Jannotta F, Liotta LA: Loss of basement membrane components by invasive tumors but not their benign counterparts. *Lab Invest* 1983, 49:140-148
- Charpin C, Lissitzky JC, Jacquemier J, Lavaut MN, Kopp F, Pourreau-Schneider N, Martin PM, Toga M: Immunohistochemical detection of laminin in 98 human breast carcinomas: A light and electron microscopic study. *Hum Pathol* 1986, 17:335-365
- Tsubara A, Shikata N, Inui T, Morii S, Hatano T, Oikawa T, Matsuzawa A: Immunohistochemical localization of myoepithelial cells and basement membrane in normal, benign, and



- malignant human breast lesions. *Virchows Arch [A]* 1988, 413:133-139
25. Barsky SH, Togo S, Garbisa S, Liotta LA: Type IV collagenase immunoreactivity in invasive breast carcinoma. *Lancet* 1983, i:296-297
  26. Hsu SM, Raine L, Fanger H: The use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase technique: A comparison between ABC and unlabeled antibody (PAP) procedures. *J Histochem Cytochem* 1981, 29:577-580
  27. Wellings SR, Jensen HM, Marcum RG: An atlas of subgross pathology of the human breast with special reference to possible precancerous lesions. *J Natl Cancer Inst* 1975, 55: 231-273
  28. Biswas C: Tumor cell stimulation of collagenase production by fibroblasts. *Biochem Biophys Res Commun* 1982, 109: 1026-1034
  29. Woolley DE, Glanville RW, Roberts DR, Evanson JM: Purification, characterization and inhibition of human skin collagenase. *Biochem J* 1978, 169:265-276
  30. Liotta LA: Gene products which play a role in cancer invasion and metastasis. *Breast Cancer Res Treat* 1988, 11: 113-124
  31. Garbisa S, Pozzatti R, Muschel RJ, Saffiotti U, Ballin M, Goldfarb RH, Khoury G, Liotta LA: Secretion of type IV collagenolytic protease and metastatic phenotype: Induction by transfection with c-Ha-ras but not c-Ha-ras plus Ad2-Ela. *Cancer Res* 1987, 47:1523-1528
  32. Bauer EA, Gordon JM, Reddick ME, Eisen AZ: Quantitation and immunocytochemical localization of human skin collagenase in basal cell carcinoma. *J Invest Dermatol* 1977, 69: 363-367
  33. Bauer EA, Uitto J, Walters RC, Eisen AZ: Enhanced collagenase production by fibroblasts derived from human basal cell carcinomas. *Cancer Res* 1979, 39:4594-4599
  34. Biswas C: Tumor cell stimulation of fibroblast collagenase production: Membrane association of the tumor cell stimulator. *Fed Proc* 1985, 44:1337
  35. Martinez-Hernandez A, Fink LM, Pierce GB: Removal of basement membrane collagen in the involuting breast. *Lab Invest* 1976, 34:455-462
  36. Wicha MS, Liotta LA, Vonderhaar BK, Kidwell WR: Effects of inhibition of basement membrane collagen deposition on rat mammary gland development. *Dev Biol* 1980, 80:253-266
  37. Ellis RA: Fine structure of the myoepithelium of the eccrine sweat glands of man. *J Cell Biol* 1965, 27:551-563
  38. Hamperl H: The myoethelia (myoepithelial cells). Normal state; regressive changes; hyperplasia; tumors. *Curr Top Pathol* 1970, 53:161-220
  39. Liotta LA, Wicha MS, Foidart JM, Rennard SI, Garbisa S, Kidwell WR: Hormonal requirements for basement membrane collagen deposition by cultured rat mammary epithelium. *Lab Invest* 1979, 41:511-518
  40. Warburton MJ, Ormerod EJ, Monaghan P, Ferns S, Rudland PS: Characterization of a myoepithelial cell line derived from a neonatal rat mammary gland. *J Cell Biol* 1981, 91:827-836
  41. Warburton MJ, Mitchell D, Ormerod J, Rudland PS: Distribution of myoepithelial cells and basement membrane proteins in the resting, pregnant, lactating, and involuting rat mammary gland. *J Histochem Cytochem* 1982, 30:667-676
  42. Nagle RB, Bocker W, Davis JR, Heid HW, Kauffmann M, Lucas DO, Jarash ED: Characterization of breast carcinomas by two monoclonal antibodies distinguishing myoepithelial from luminal epithelial cells. *J Histochem Cytochem* 1986, 34:869-881
  43. Jarash E-D, Nagle RB, Kaufman M, Maurer C, Bocker WJ: Differential diagnosis of benign epithelial proliferations and carcinomas of the breast using antibodies to cytokeratins. *Hum Pathol* 1988, 19:276-289

## Acknowledgment

The authors thank Dr. Ronald Neumann for providing the laboratory and technical resources to perform the immunohistochemical study.